

The Conley Index for Decompositions of Isolated Invariant Sets ¹

by

Andrzej Szymczak (Gdańsk) ²

Abstract

Let f be a continuous map of a locally compact metric space X into itself. Suppose that S is an isolated invariant set with respect to f being a disjoint union of a fixed finite number of compact sets. We define an index of Conley type for isolated invariant sets admitting such a decomposition and prove some of its properties, which appear to be similar to that of the ordinary Conley index for maps. Our index takes into account the existence of the decomposition of S and therefore carries more information about the structure of the invariant set. In particular, it seems to be a more accurate tool for the detection of periodic trajectories and chaos of the Smale horseshoe type than the ordinary Conley index.

0. Introduction

The Conley index has become an important tool in the study of the qualitative behaviour of dynamical systems, with both discrete and continuous time. The results concerning attractor-repeller decompositions ([1], [15], [18]), the connection matrix theory ([3], [4], [5]) as well as recent papers by Ch.McCord, K.Mischaikow and M.Mrozek [7] and the latter two authors [9] (see also [20]) show that the Conley index reflects the structure of an isolated invariant set. In this paper we are mainly interested in the Conley index as a tool for the detection of chaos and periodic orbits. Comparing the results of [9] and [20] with the criteria for chaos based on the fixed point index in [19] or [23] shows that the ones based on the Conley index

¹1991 *Mathematics Subject Classification* : 54H20,34C35

²Supported by UG grant BW 5100-5-0092-4

are, in some sense, weak. They only guarantee that some iteration of the map restricted to the isolated invariant set is semiconjugate with the shift map. Thus, they provide information about the dynamics of some iteration of the map rather than the map itself. The information about the number of periodic orbits is also not as accurate as that provided by the methods based on the fixed point index. The aim of this paper is to define an index of Conley type which fills this gap. Our index is defined for a decomposition of an isolated invariant set into a fixed number of disjoint compact sets. The knowledge of the decomposition allows to equip the index with an additional structure, which carries more information than the ordinary Conley index. The main potential application of our index is for the detection of chaos. The Conley index for decompositions can also be used to state topological analogues of some results in the theory of smooth dynamical systems (e.g. the Poincare-Birkhoff theorem). This topic will be discussed in a separate paper.

1. Preliminaries

By Z^+ and R we shall denote the sets of nonnegative integer and real numbers (respectively). If X is a metric space and $Q = (Q_1, Q_0)$ is a pair of its compact subsets then by Q_1/Q_0 we denote the pointed space resulting from Q_1 when the points of Q_0 are identified to a single distinguished point, denoted by $[Q_0]$. $\mathcal{H}top$, \mathcal{M} and \mathcal{M}_G will stand for the homotopy category of pointed topological spaces, the category of modules and the category of graded modules over a fixed ring with unity Ξ . For a basepoint preserving map g its homotopy class will also be denoted by g . This should not cause misunderstanding. For an object O in a category \mathcal{K} by $[O]$ we shall denote the class of all objects in \mathcal{K} isomorphic to O . A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ induces the map sending an isomorphism class $[O]$ into $[F(O)]$ for each object $O \in Ob(\mathcal{K})$.

We shall denote this map by the same letter F .

Let us recall the basic concepts of the Conley index theory now. Our presentation is based mainly on [21] and [20] (see also [11], [13], [16], [17]). We begin with the definition of the category of objects equipped with a morphism over a given category \mathcal{K} , denoted by \mathcal{K}_m . Put

$$Ob(\mathcal{K}_m) = \{(X, \alpha) : X \in Ob(\mathcal{K}) \text{ and } \alpha \in Mor_{\mathcal{K}}(X, X)\}$$

and

$$Mor_{\mathcal{K}_m}((X, \alpha), (X', \alpha')) = M((X, \alpha), (X', \alpha')) / \equiv$$

where

$$M((X, \alpha), (X', \alpha')) = \{\beta \in Mor_{\mathcal{K}}(X, X') : \beta \circ \alpha = \alpha' \circ \beta\} \times Z^+$$

and \equiv is the equivalence relation in the above set defined by

$$(\beta, n) \equiv (\bar{\beta}, \bar{n}) \iff \exists_{k \in Z^+} \beta \circ \alpha^{\bar{n}+k} = \bar{\beta} \circ \alpha^{n+k}.$$

The morphism represented by $(\beta, n) \in M((X, \alpha), (X', \alpha'))$ will be denoted by $[\beta, n]$. The composition of morphisms in \mathcal{K}_m is defined by

$$[\beta', n'] \circ [\beta, n] = [\beta' \circ \beta, n' + n].$$

Given a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ one can define the induced functor $F_m : \mathcal{K}_m \rightarrow \mathcal{L}_m$ in the following way. For an object (X, α) and a morphism $[\beta, n]$ in \mathcal{K}_m we put

$$F_m(X, \alpha) = (F(X), F(\alpha)),$$

$$F_m([\beta, n]) = [F(\beta), n].$$

In the sequel we shall use the notation $[X, \alpha]$ for the class of all objects in \mathcal{K}_m isomorphic to an object (X, α) .

In each of the categories $\mathcal{H}top$, \mathcal{M} , \mathcal{M}_G for each object X there exists the zero morphism of X into itself (i.e. the homotopy class of the constant map or the zero homomorphism, according to the case). We shall denote this morphism by 0 . In the same way we shall denote the trivial isomorphism classes in the categories of objects equipped with a morphism over each of the three categories, i.e. we put $0 = [C, 0]$ where C is any pointed space or (graded) Ξ -module. This class is independent on the choice of C . This notation is ambiguous, but it will always be clear from context what is meant by 0 . We note that $[X, \alpha] = 0$ if and only if $\alpha^n = 0$ for some $n \in \mathbb{Z}^+$.

For the rest of this section, fix a locally compact metric space X and a continuous map f of this space into itself. Let S be an isolated invariant set with respect to f . A pair $Q = (Q_1, Q_0)$ of compact subsets of X is called an index pair for S with respect to f if and only if $S = Inv_f cl(Q_1 \setminus Q_0) \subset int(Q_1 \setminus Q_0)$, Q_0 is positively invariant in Q_1 (i.e. $f(Q_0) \cap Q_1 \subset Q_0$) and Q_0 is an exit set for Q_1 (which means that $f(Q_1 \setminus Q_0) \subset Q_1$). For such Q , f induces the continuous map $f_Q : Q_1/Q_0 \rightarrow Q_1/Q_0$ which will be called the index map. The (homotopy) Conley index of S , denoted by $h(S, f, X)$ is defined as the class of all objects in $\mathcal{H}top_m$ isomorphic to $(Q_1/Q_0, f_Q)$. We define the cohomological and the q -dimensional cohomological Conley indices by

$$h^*(S, f, X) = (H^*)_m(h(S, f, X))$$

and

$$h^q(S, f, X) = (H^q)_m(h(S, f, X))$$

where $H^* : \mathcal{H}top \rightarrow \mathcal{M}_G$ is a fixed cohomology functor with coefficients in Ξ .

To the end of this section, let Ξ be the field of rational numbers. Then, \mathcal{M} and \mathcal{M}_G are the categories of vector spaces and graded vector spaces over

this field. An object (V, φ) in \mathcal{M}_m is called of finite asymptotic dimension (cf [20], definition 2.1 and proposition 2.2) if and only if there exists an object (W, ψ) with W finite dimensional, isomorphic to (V, φ) . In this case, we define the trace of (V, φ) , denoted by $tr(V, \varphi)$ as the ordinary trace of ψ . Using the methods of [20] (see theorem 1.1, definition 4.1 and remark 4.1) one proves easily that it is independent on the choice of (W, ψ) . Now, let (V^*, φ^*) be an object in $(\mathcal{M}_G)_m$. It is said to be of finite type if and only if there exists an object (W^*, ψ^*) , with W^* of finite type, isomorphic to (V^*, φ^*) . In this case, we define the Lefschetz number of (V^*, φ^*) , denoted by $\Lambda(V^*, \varphi^*)$, as the ordinary Lefschetz number of ψ^* . Clearly,

$$\Lambda(V^*, \varphi^*) = \sum_{q=-\infty}^{\infty} (-1)^q tr(V^q, \varphi^q).$$

An isomorphism class I of objects in $(\mathcal{M}_G)_m$ is said to be of finite type if it admits a representative being of finite type. In this case, all of its representatives are of finite type and they have the same Lefschetz number, which we call the Lefschetz number of I and denote by $\Lambda(I)$. The following theorem is taken from [20] (see lemma 5.2). For related results, see [10],[12],[14].

Theorem 1.1 *If X is an Euclidean neighborhood retract (ENR), then the cohomological Conley index of any isolated invariant set with respect to f is of finite type. If the Lefschetz number of the Conley index of an isolated invariant set S is nonzero then f has a fixed point in S .*

2. Categorical constructions

There are a lot of Conley-type indices for isolated invariant sets (see [1], [11], [13], [17], [18], [21]), but all of them take the form of an isomorphism class of objects in a certain category. In the classical, continuous-time case, the homotopy category of pointed topological spaces is used and therefore

the Conley index is simply a homotopy class of a pointed space. In the discrete-time case the situation is much more complicated: in order to give a good definition one has to use more sophisticated categorical constructions. The shape category ([13], [17]), the Leray functor ([11], [13]), the direct and inverse limit functors ([13]) and the category of objects equipped with a morphism ([21]) can serve as examples here. Below we define a generalization of the latter concept, which is suitable for the definition of the Conley index for decompositions of isolated invariant sets.

Let \mathcal{K} be a category and A a finite set. In the sequel, we shall often deal with finite sequences of elements of A . By $*$ we shall denote the concatenation operation on the set of such sequences, defined by

$$(Z_1, Z_2, \dots, Z_n) * (Z'_1, Z'_2, \dots, Z'_m) = (Z_1, Z_2, \dots, Z_n, Z'_1, Z'_2, \dots, Z'_m) \in A^{n+m}$$

for all $(Z_1, Z_2, \dots, Z_n) \in A^n$ and $(Z'_1, Z'_2, \dots, Z'_m) \in A^m$. For \bar{Z} being a sequence of members of A by \bar{Z}^k we shall denote the result of the $*$ operation performed on k copies of \bar{Z} . By $\iota(\bar{Z})$ we shall denote the sequence \bar{Z} with the entries in the reversed order.

Let us define the category $\mathcal{K}_{(A)}$ now. For $n \in \mathbb{Z}^+$ and $X, X' \in \text{Ob}(\mathcal{K})$ put:

$$\begin{aligned} \text{Ob}(\mathcal{K}_{(A)}) &= \text{Ob}(\mathcal{K}), \\ \text{Mor}_{\mathcal{K}_{(A)}}^n(X, X') &= (\text{Mor}_{\mathcal{K}}(X, X'))^{A^n}, \\ \text{Mor}_{\mathcal{K}_{(A)}}(X, X') &= \bigcup_{n \in \mathbb{Z}^+} \text{Mor}_{\mathcal{K}_{(A)}}^n(X, X'). \end{aligned}$$

The composition of morphisms

$$\alpha = \{\alpha^{\bar{Z}}\}_{\bar{Z} \in A^n} \in \text{Mor}_{\mathcal{K}_{(A)}}^n(X, X')$$

and

$$\alpha' = \{\alpha'^{\bar{Z}}\}_{\bar{Z} \in A^m} \in \text{Mor}_{\mathcal{K}_{(A)}}^m(X', X'')$$

is defined as follows:

$$\alpha' \circ \alpha = \{(\alpha' \circ \alpha)^{\bar{Z}}\}_{\bar{Z} \in A^{m+n}} \in Mor_{\mathcal{K}_{(A)}}^{m+n}(X, X'')$$

where

$$(\alpha' \circ \alpha)^{\bar{Y}' * \bar{Y}} = \alpha'^{\bar{Y}'} \circ \alpha^{\bar{Y}}.$$

for all $\bar{Y} \in A^n$ and $\bar{Y}' \in A^m$. Since A^0 consists of exactly one element, we shall identify $Mor_{\mathcal{K}_{(A)}}^0(X, X')$ with $Mor_{\mathcal{K}}(X, X')$ in the obvious way. Similarly, there is an obvious bijection of $Mor_{\mathcal{K}}^1(X, X')$ onto $(Mor_{\mathcal{K}}(X, X'))^A$. Therefore, we treat morphisms in this set as families of morphisms of X into X' in \mathcal{K} , indexed by members of A . It is straightforward to verify that $\mathcal{K}_{(A)}$ is indeed a category. Notice that its identity morphism over an object X is $id_X \in Mor_{\mathcal{K}_{(A)}}^0(X, X)$.

The $\mathcal{K}_{(A)}$ category is only an intermediate step in the definition of the $\mathcal{K}_{[A]}$ category, which we are going to use in the definition of the Conley index for decompositions of isolated invariant sets. Put

$$Ob(\mathcal{K}_{[A]}) = \{(X, \alpha) : X \in Ob(\mathcal{K}_{(A)}) = Ob(\mathcal{K}), \alpha \in Mor_{\mathcal{K}_{(A)}}^1(X, X)\}.$$

In order to define morphisms in $\mathcal{K}_{[A]}$, for objects (X, α) and (X', α') put

$$M((X, \alpha), (X', \alpha')) = \{(\beta, n) : \beta \in Mor_{\mathcal{K}_{(A)}}^n(X, X'), n \in \mathbb{Z}^+, \beta \circ \alpha = \alpha' \circ \beta\}.$$

In this set we introduce the equivalence relation \equiv in the following way.

$$(\beta, n) \equiv (\bar{\beta}, \bar{n}) \iff \exists_{k \in \mathbb{Z}^+} \beta \circ \alpha^{\bar{n}+k} = \bar{\beta} \circ \alpha^{n+k} \quad (\text{in } \mathcal{K}_{(A)}).$$

Now, define

$$Mor_{\mathcal{K}_{[A]}}((X, \alpha), (X', \alpha')) = M((X, \alpha), (X', \alpha')) / \equiv.$$

The morphism represented by (β, n) will be denoted by $[\beta, n]$. The composition of morphisms

$$[\beta, n] : (X, \alpha) \longrightarrow (X', \alpha')$$

and

$$[\beta', n'] : (X', \alpha') \longrightarrow (X'', \alpha'')$$

is defined as follows

$$[\beta', n'] \circ [\beta, n] = [\beta' \circ \beta, n' + n].$$

One can easily verify that this definition is correct, i.e. independent on the choice of representatives for $[\beta, n]$ and $[\beta', n']$ and that $\mathcal{K}_{[A]}$ is indeed a category. Note that the identity morphism over (X, α) is $[id_X, 0]$.

Proposition 2.1 *For each $[\beta, n] \in Mor_{\mathcal{K}_{[A]}}((X, \alpha), (X', \alpha'))$ and $k \in Z^+$*

$$[\beta, n] = [\alpha'^k \circ \beta, n + k] = [\beta \circ \alpha^k, n + k].$$

Proof. This follows immediately from the definition of \equiv . \square

The Conley index for decompositions of isolated invariant sets will 'contain' information about ordinary Conley indices of some sets which are important for understanding of the dynamics of the map. Below we give the definition of functors which will enable us to extract this information.

Let k be a positive integer and $\bar{Y} = (Y_1, Y_2, \dots, Y_k) \in A^k$. The functor $\mathcal{P}_{\bar{Y}} : \mathcal{K}_{[A]} \longrightarrow \mathcal{K}_m$ is defined as follows. For an object (X, α) in $\mathcal{K}_{[A]}$ with $\alpha = \{\alpha^Z\}_{Z \in A}$ put

$$\mathcal{P}_{\bar{Y}}(X, \alpha) = (X, \alpha^{Y_1} \circ \alpha^{Y_2} \circ \dots \circ \alpha^{Y_k}).$$

Now, let $[\beta, n]$ be a morphism of (X, α) into (X', α') . By proposition 2.1, without the loss of generality we can assume that $n = mk$ for some $m \in Z^+$. Suppose that $\beta = \{\beta^{\bar{Z}}\}_{\bar{Z} \in A^n}$. Put

$$\mathcal{P}_{\bar{Y}}([\beta, n]) = [\beta^{\bar{Y}^m}, m].$$

A routine check that $\mathcal{P}_{\bar{Y}}$ is a well-defined functor is left to the reader.

Remark 2.1. An important property of the construction given above is the naturality with respect to functors. Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a functor. Then we have the induced functors $F_{(A)} : \mathcal{K}_{(A)} \rightarrow \mathcal{L}_{(A)}$ and $F_{[A]} : \mathcal{K}_{[A]} \rightarrow \mathcal{L}_{[A]}$ defined as follows.

$$F_{(A)}(X) = F(X),$$

$$F_{(A)}(\{\alpha^{\bar{Z}}\}_{\bar{Z} \in A^k}) = \begin{cases} \{F(\alpha^{\bar{Z}})\}_{\bar{Z} \in A^k} & \text{if } F \text{ is covariant} \\ \{F(\alpha^{\iota(\bar{Z})})\}_{\bar{Z} \in A^k} & \text{if } F \text{ is contravariant} \end{cases}$$

for all objects X and morphisms $\{\alpha^{\bar{Z}}\}_{\bar{Z} \in A^k}$ in $\mathcal{K}_{(A)}$ and

$$F_{[A]}(X, \alpha) = (F_{(A)}(X), F_{(A)}(\alpha)),$$

$$F_{[A]}([\beta, n]) = [F_{(A)}(\beta), n]$$

for all objects (X, α) and morphisms $[\beta, n]$ in $\mathcal{K}_{[A]}$. Furthermore, the following diagram of categories and functors commutes for each $\bar{Y} \in A^k$

$$\begin{array}{ccc} \mathcal{K}_{[A]} & \xrightarrow{F_{[A]}} & \mathcal{L}_{[A]} \\ \downarrow \mathcal{P}_{\bar{Y}} & & \downarrow \mathcal{P}_{\iota_F(\bar{Y})} \\ \mathcal{K}_m & \xrightarrow{F_m} & \mathcal{L}_m \end{array}$$

where

$$\iota_F(\bar{Y}) = \begin{cases} \iota(\bar{Y}) & \text{if } F \text{ is contravariant} \\ \bar{Y} & \text{if } F \text{ is covariant} \end{cases}.$$

As in the case of categories equipped with a morphism, by 0 we shall denote the isomorphism classes of the trivial (zero) objects in $\mathcal{Htop}_{[A]}$, $(\mathcal{M}_G)_{[A]}$ and $\mathcal{M}_{[A]}$, defined in the obvious way.

3. The index

This section contains the basic definitions of the Conley index theory for decompositions of isolated invariant sets. In what follows, B, X and f will denote a fixed finite set, a locally compact metric space and a continuous map of X into itself. We shall make use of the categories of $\mathcal{K}_{[A]}$ type with $A = 2^B$. If $\{N_b\}$ is a family of sets indexed by members of B then for each set $Z \subset B$ by N_Z we shall denote the union of N_b over $b \in Z$.

Definition 3.1 *Let N be a compact subset of X . A family $\{N_b\} = \{N_b\}_{b \in B}$ of pairwise disjoint compact sets is called a decomposition of N if $N = \bigcup_{b \in B} N_b$.*

From now to the end of this section, by S we shall denote a fixed isolated invariant set with respect to f and by $\{S_b\}$ its decomposition.

Definition 3.2 *An index pair $Q = (Q_1, Q_0)$ for S is said to be compatible with the decomposition $\{S_b\}$ of S if and only if there exists $\{D_b\}$ a decomposition of $cl(Q_1 \setminus Q_0)$ such that $S_b = S \cap D_b$ for each $b \in B$.*

Let us emphasise that, in general, the decomposition $\{D_b\}$ is not uniquely determined by Q and $\{S_b\}$.

Definition 3.3 *Let $Q = (Q_1, Q_0)$ be an index pair for S compatible with the decomposition $\{S_b\}$ of S and $\{D_b\}$ be a corresponding decomposition of $cl(Q_1 \setminus Q_0)$. Then, for any $Z \in A$ we can define the continuous map*

$$r^Z = r^Z_{(Q, \{D_b\})} : Q_1/Q_0 \longrightarrow Q_1/Q_0$$

by the following formula

$$r^Z([x]) = \begin{cases} [x] & \text{if } x \in D_Z \\ [Q_0] & \text{otherwise} \end{cases} .$$

The index object, denoted by $I(Q, \{D_b\}, f)$, is the object in $\mathcal{Htop}_{[A]}$ given by

$$I(Q, \{D_b\}, f) = (Q_1/Q_0, \{f^Z\}_{Z \in A})$$

where $f^Z = f^Z_{(Q, \{D_b\})} = f_Q \circ r^Z$ (recall that f_Q is the index map).

In order to simplify the notation, we shall often write briefly $I(Q, \{D_b\})$ instead of $I(Q, \{D_b\}, f)$ whenever the map f is clear from context. For each $Z \in A$ and $x \in Q_1$ we have the following formula

$$f^Z([x]) = \begin{cases} [f(x)] & \text{if } x \in D_Z \cap (Q_1 \setminus Q_0) \\ [Q_0] & \text{otherwise} \end{cases}$$

For further reference, let us note the following formula for compositions of the maps f^Z . Let $\bar{Z} = (Z_0, Z_1, \dots, Z_{T-1}) \in A^T$. For all $x \in Q_1$,

$$\begin{aligned} f^{Z_{T-1}} \circ f^{Z_{T-2}} \circ \dots \circ f^{Z_0}([x]) &= \\ &= \begin{cases} [f^T(x)] & \text{if } f^i(x) \in D_{Z_i} \cap (Q_1 \setminus Q_0) \\ & \text{for each } i \in \{0, 1, \dots, T-1\} \\ [Q_0] & \text{otherwise} \end{cases} \end{aligned} \quad (3.1)$$

Let N be a compact neighborhood of S admitting a decomposition $\{N_b\}$ such that $S_b = N_b \cap S$ for each $b \in B$. By the existence theorems for index pairs (see [6], [10], [11], [13], [16]) there is an index pair $Q = (Q_1, Q_0)$ for S with Q_1 contained in N (Q may be even assumed regular in the sense of [10] or [20]). Obviously, such an index pair is compatible with the decomposition $\{S_b\}$. We have proved the following

Proposition 3.1 *There exist index pairs for S compatible with the decomposition $\{S_b\}$, arbitrarily close to S .*

The following theorem is of fundamental importance in our construction.

Theorem 3.1 *If $Q = (Q_1, Q_0)$ and $\bar{Q} = (\bar{Q}_1, \bar{Q}_0)$ are index pairs for S , compatible with the decomposition $\{S_b\}$ of S and $\{D_b\}$ and $\{\bar{D}_b\}$ are decom-*

positions of $cl(Q_1 \setminus Q_0)$ and $cl(\bar{Q}_1 \setminus \bar{Q}_0)$ satisfying the conditions of definition 3.2 then the index objects $I(Q, \{D_b\})$ and $I(\bar{Q}, \{\bar{D}_b\})$ are isomorphic in $\mathcal{H}top_{[A]}$.

Proof. We proceed in several steps.

Step 1. There exists $T \in Z^+$ such that the following two implications hold for each sequence $(Z_0, Z_1, \dots, Z_{2T-1})$ of members of A and $x \in X$.

$$(\forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in \bar{D}_{Z_i}) \Rightarrow f^T(x) \in D_{Z_T} \cap (Q_1 \setminus Q_0) \quad (3.2)$$

and

$$(\forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in D_{Z_i}) \Rightarrow f^T(x) \in \bar{D}_{Z_T} \cap (\bar{Q}_1 \setminus \bar{Q}_0). \quad (3.3)$$

For the proof, notice that the set U given by

$$U = \bigcup_{b \in B} (D_b \cap \bar{D}_b \cap (Q_1 \setminus Q_0) \cap (\bar{Q}_1 \setminus \bar{Q}_0))$$

is a neighborhood of S . As a consequence of lemma 4.2 in [21] (cf also lemma 6.2 in [17]) we obtain the existence of a nonnegative integer T such that the following two implications holds for each $x \in X$.

$$\begin{aligned} (\forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in cl(Q_1 \setminus Q_0)) &\Rightarrow f^T(x) \in U, \\ (\forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in cl(\bar{Q}_1 \setminus \bar{Q}_0)) &\Rightarrow f^T(x) \in U \end{aligned} \quad (3.4)$$

Now, suppose that $\forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in D_{Z_i}$. Then, by (3.4), $f^T(x) \in U$.

Since simultaneously $f^T(x) \in D_{Z_T}$,

$$f^T(x) \in U \cap D_{Z_T} \subset \bar{D}_{Z_T} \cap (\bar{Q}_1 \setminus \bar{Q}_0).$$

We have proved that the implication (3.3) holds. In a similar way one proves that so does (3.2).

Step 2. Let $T \in Z^+$ be such that (3.2) and (3.3) hold. For a sequence $\bar{Z} = (Z_1, Z_2, \dots, Z_{3T})$ of members of A we define the function

$$f^{\bar{Z}} = f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})}^{\bar{Z}} : Q_1/Q_0 \longrightarrow \bar{Q}_1/\bar{Q}_0$$

by the following formula

$$f^{\bar{Z}}([x]) = \begin{cases} [f^{3T}(x)] & \text{if } \forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in D_{Z_{3T-i}} \cap (Q_1 \setminus Q_0) \\ & \text{and } f^{T+i}(x) \in \bar{D}_{Z_{2T-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0) \\ [\bar{Q}_0] & \text{otherwise} \end{cases}.$$

Our task is to prove the continuity of $f^{\bar{Z}}$.

The proof goes along the lines of other continuity proofs in the Conley index theory (cf [17], [18], [21]). Let $f_0^{\bar{Z}} : Q_1 \rightarrow \bar{Q}_1/\bar{Q}_0$ be defined by the same formula as $f^{\bar{Z}}$. Clearly, it is enough to show that $f_0^{\bar{Z}}$ is continuous. Put

$$\begin{aligned} O_1 &= \{x \in Q_1 : \forall_{i \in \{0, \dots, 2T-1\}} f^i(x) \in D_{Z_{3T-i}} \cap (Q_1 \setminus Q_0) \\ &\quad \text{and } f^{T+i}(x) \in \bar{D}_{Z_{2T-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0)\}, \\ O_2 &= \{x \in Q_1 : \exists_{i \in \{0, \dots, 2T-1\}} f^i(x) \notin D_{Z_{3T-i}} \text{ or } f^{T+i}(x) \notin \bar{D}_{Z_{2T-i}}\}. \end{aligned}$$

Clearly, O_2 is open in Q_1 and $f_0^{\bar{Z}}$ is constant on O_2 and therefore continuous at each point of this set. Since $f_0^{\bar{Z}}(x) = [f^{3T}(x)]$ for each $x \in O_1$, in order to prove the continuity of $f_0^{\bar{Z}}$ at each point of O_1 it is enough to show that this set is open in Q_1 . Take $x \in O_1$. There exists U an open neighborhood of x in X such that

$$f^i(U) \cap (Q_0 \cup D_{B \setminus Z_{3T-i}}) = \emptyset = f^{T+i}(U) \cap (\bar{Q}_0 \cup \bar{D}_{B \setminus Z_{2T-i}}) \quad (3.5)$$

for each $i \in \{0, 1, \dots, 2T-1\}$. Let us show that $U \cap Q_1 \subset O_1$. Let $y \in U \cap Q_1$. Our assumptions about U imply $y \in Q_1 \setminus Q_0$ and $f^i(y) \notin Q_0$ for each $i \in \{0, 1, \dots, 2T-1\}$. Since Q_0 is an exit set for Q_1 , $f^i(y) \in Q_1 \setminus Q_0$. By (3.5), $f^i(y) \in (Q_1 \setminus Q_0) \cap D_{Z_{3T-i}}$. Hence, by (3.3),

$$f^T(y) \in \bar{D}_{Z_{2T}} \cap (\bar{Q}_1 \setminus \bar{Q}_0) \subset \bar{Q}_1 \setminus \bar{Q}_0.$$

By the previous argument,

$$f^{T+i}(y) \in \bar{D}_{Z_{2T-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0) \quad \text{for all } i \in \{0, 1, \dots, 2T-1\}$$

so that $y \in O_1$.

We conclude that, in order to prove the continuity of $f_0^{\bar{Z}}$, it is enough to show that it is continuous at each point of $Q_1 \setminus (O_1 \cup O_2)$. Let x be a member of this set. Then, in particular,

$$\forall_{i \in \{0, \dots, 2T-1\}} \quad f^i(x) \in D_{Z_{3T-i}} \text{ and } f^{T+i}(x) \in \bar{D}_{Z_{2T-i}} \quad (3.6)$$

and $f_0^{\bar{Z}}(x) = [\bar{Q}_0]$. By (3.2) applied for x replaced with $f^T(x)$, $f^{2T}(x) \in D_{Z_T} \cap (Q_1 \setminus Q_0)$. Since $f^i(x) \in D_{Z_{3T-i}} \subset Q_1$ for each $i \in \{0, 1, \dots, 2T-1\}$, positive invariance of Q_0 in Q_1 implies $f^i(x) \in D_{Z_{3T-i}} \cap (Q_1 \setminus Q_0)$. Thus, since $x \notin O_1$, for some $j \in \{0, 1, \dots, 2T-1\}$ we must have $f^{T+j}(x) \notin \bar{D}_{Z_{2T-j}} \cap (\bar{Q}_1 \setminus \bar{Q}_0)$. By (3.6), $f^{T+j}(x) \in \bar{Q}_0$. By (3.6) and positive invariance of \bar{Q}_0 in \bar{Q}_1 , $f^{3T-1}(x) \in \bar{Q}_0$. Let V' be an open neighborhood of $[\bar{Q}_0]$ in \bar{Q}_1/\bar{Q}_0 . By π we shall denote the projection map of \bar{Q}_1 into \bar{Q}_1/\bar{Q}_0 . Put $M = \pi^{-1}((\bar{Q}_1/\bar{Q}_0) \setminus V')$. Clearly, M is a compact subset of $\bar{Q}_1 \setminus \bar{Q}_0$. By positive invariance of \bar{Q}_0 in \bar{Q}_1 , $f^{3T}(x) \notin M$. Let V be an open neighborhood of x in Q_1 such that $f^{3T}(V) \cap M = \emptyset$. Notice that for all $y \in V$, $f_0^{\bar{Z}}(y)$ is equal to either $[\bar{Q}_0]$ or $[f^{3T}(y)]$ (the second possibility can occur only if $f^{3T}(y) \in \bar{Q}_1$). Hence, $f_0^{\bar{Z}}(y) \in V'$. In this way we have proved that $f_0^{\bar{Z}}$ is continuous at x .

Step 3. If $(Z_1, Z_2, \dots, Z_{3T+1}) \in A^{3T+1}$ then

$$f^{(Z_1, \dots, Z_{3T})} \circ f_{(Q, \{D_b\})}^{Z_{3T+1}} = f_{(\bar{Q}, \{\bar{D}_b\})}^{Z_1} \circ f^{(Z_2, \dots, Z_{3T+1})}.$$

Therefore, we have the following morphism in $Htop_{[A]}$.

$$f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})} = [\{f^{\bar{Z}}\}_{\bar{Z} \in A^{3T}, 3T}] : I(Q, \{D_b\}) \longrightarrow I(\bar{Q}, \{\bar{D}_b\}).$$

To prove this, consider the following two conditions:

$$\begin{aligned} \forall_{i \in \{0, \dots, 2T\}} \quad f^i(x) &\in D_{Z_{3T+1-i}} \cap (Q_1 \setminus Q_0) && \text{and} \\ \forall_{i \in \{1, \dots, 2T\}} \quad f^{T+i}(x) &\in \bar{D}_{Z_{2T+1-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0) \end{aligned} \quad (3.7)$$

and

$$\begin{aligned} \forall_{i \in \{0, \dots, 2T-1\}} \quad f^i(x) &\in D_{Z_{3T+1-i}} \cap (Q_1 \setminus Q_0) \quad \text{and} \\ \forall_{i \in \{0, \dots, 2T\}} \quad f^{T+i}(x) &\in \bar{D}_{Z_{2T+1-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0). \end{aligned} \quad (3.8)$$

Notice that if for some $x \in X$ (3.7) holds then applying (3.3) gives $f^T(x) \in \bar{D}_{Z_{2T+1}} \cap (\bar{Q}_1 \setminus \bar{Q}_0)$. Therefore, (3.8) holds. We have proved that (3.7) \Rightarrow (3.8). Since the reverse implication can be proved in a similar way using (3.2) with x replaced with $f^T(x)$, (3.7) and (3.8) are equivalent. To finish the proof apply the formulas for f^Z and $f^{\bar{Z}}$ to obtain that, for all $x \in Q_1$,

$$f^{(Z_1, \dots, Z_{3T})} \circ f_{(Q, \{D_b\})}^{Z_{3T+1}}([x]) = \begin{cases} [f^{3T+1}(x)] & \text{if (3.7) holds} \\ [Q_0] & \text{otherwise} \end{cases}$$

and

$$f_{(\bar{Q}, \{\bar{D}_b\})}^{Z_1} \circ f^{(Z_2, \dots, Z_{3T+1})} = \begin{cases} [f^{3T+1}(x)] & \text{if (3.8) holds} \\ [\bar{Q}_0] & \text{otherwise} \end{cases}.$$

Step 4. The morphism $f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})}$ defined in step 3 is an isomorphism in $Htop[A]$.

Notice that steps 1 through 3 remain valid if we replace Q by \bar{Q} and vice versa. Hence we have the morphism

$$\begin{aligned} f_{(\bar{Q}, \{\bar{D}_b\}), (Q, \{D_b\})} &= \\ &= \left[\{ f_{(\bar{Q}, \{\bar{D}_b\}), (Q, \{D_b\})}^{\bar{Z}} \}_{\bar{Z} \in A^{3T}, 3T} \right] : I(\bar{Q}, \{\bar{D}_b\}) \longrightarrow I(Q, \{D_b\}). \end{aligned}$$

Now, consider the composition

$$g = f_{(\bar{Q}, \{\bar{D}_b\}), (Q, \{D_b\})}^{\bar{Y}} \circ f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})}^{\bar{Z}}$$

for given $\bar{Z} = (Z_1, Z_2, \dots, Z_{3T}) \in A^{3T}$ and $\bar{Y} = (Y_1, Y_2, \dots, Y_{3T}) \in A^{3T}$.

Using the formula used to define the maps in step 2 we get

$$\begin{aligned} g([x]) &= \begin{cases} [f^{6T}(x)] & \text{if the condition (3.9) given below holds} \\ [Q_0] & \text{otherwise} \end{cases}, \\ \forall_{i \in \{0, \dots, 2T-1\}} \quad f^i(x) &\in D_{Z_{3T-i}} \cap (Q_1 \setminus Q_0), \\ f^{T+i}(x) &\in \bar{D}_{Z_{2T-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0), \\ f^{3T+i}(x) &\in \bar{D}_{Y_{3T-i}} \cap (\bar{Q}_1 \setminus \bar{Q}_0) \\ \text{and} \quad f^{4T+i}(x) &\in D_{Y_{2T-i}} \cap (Q_1 \setminus Q_0). \end{aligned} \quad (3.9)$$

Using the implications (3.2) and (3.3) one can easily prove that the above condition is equivalent to the following one

$$\forall i \in \{0, \dots, 3T-1\} f^i(x) \in D_{Z_{3T-i}} \cap (Q_1 \setminus Q_0) \text{ and } f^{3T+i}(x) \in D_{Y_{3T-i}} \cap (Q_1 \setminus Q_0).$$

Hence, by the formula (3.1),

$$g = f_{(Q, \{D_b\})}^{Y_1} \circ f_{(Q, \{D_b\})}^{Y_2} \circ \dots \circ f_{(Q, \{D_b\})}^{Y_{3T}} \circ f_{(Q, \{D_b\})}^{Z_1} \circ f_{(Q, \{D_b\})}^{Z_2} \circ \dots \circ f_{(Q, \{D_b\})}^{Z_{3T}}.$$

This means that

$$\begin{aligned} & \{f_{(\bar{Q}, \{\bar{D}_b\}), (Q, \{D_b\})}^{\bar{Z}}\}_{\bar{Z} \in A^{3T}} \circ \{f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})}^{\bar{Z}}\}_{\bar{Z} \in A^{3T}} = \\ & = (\{f_{(Q, \{D_b\})}^Z\}_{Z \in A})^{6T} \end{aligned}$$

in $Htop_{(A)}$ and therefore, by proposition 2.1,

$$\begin{aligned} & f_{(\bar{Q}, \{\bar{D}_b\}), (Q, \{D_b\})} \circ f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})} = \\ & = [(\{f_{(Q, \{D_b\})}^Z\}_{Z \in A})^{6T}, 6T] = id_{I(Q, \{D_b\})}. \end{aligned}$$

In a similar way one proves that

$$f_{(Q, \{D_b\}), (\bar{Q}, \{\bar{D}_b\})} \circ f_{(\bar{Q}, \{\bar{D}_b\}), (Q, \{D_b\})} = id_{I(\bar{Q}, \{\bar{D}_b\})}. \quad \square$$

The just proved theorem justifies the following definition.

Definition 3.4 *Let $f : X \rightarrow X$ be a continuous map, S an isolated invariant set with respect to f and $\{S_b\}$ its decomposition. The Conley index of $\{S_b\}$, denoted by $h(\{S_b\}, f, X)$ is defined as the class of all objects in $Htop_{[A]}$ isomorphic to the index object $I(Q, \{D_b\})$ for any index pair Q for S compatible with the decomposition $\{S_b\}$ and any decomposition $\{D_b\}$ of $cl(Q_1 \setminus Q_0)$ such that $D_b \cap S = S_b$ for each $b \in B$. The cohomological and the q -dimensional cohomological indices of $\{S_b\}$ are defined by*

$$h^*(\{S_b\}, f, X) = (H^*)_{[A]}(h(\{S_b\}, f, X))$$

and

$$h^q(\{S_b\}, f, X) = (H^q)_{[A]}(h(\{S_b\}, f, X)),$$

respectively.

4. Properties

We begin this section with proving the continuation property of the Conley index for decompositions of isolated invariant sets.

Theorem 4.1 (Continuation property) *Suppose that X is a locally compact metric space and a continuous map $f : X \times [0, 1] \rightarrow X \times [0, 1]$ is parameter-preserving, i. e. for each $\lambda \in [0, 1]$ it satisfies $f(X \times \lambda) \subset X \times \lambda$. For each set $A \subset X \times [0, 1]$ and $\lambda \in [0, 1]$ we put $A_\lambda = \{x \in X : (x, \lambda) \in A\}$. The map $f_\lambda : X \rightarrow X$ is defined by $f(x, \lambda) = (f_\lambda(x), \lambda)$. If S is an isolated invariant set with respect to f and $\{S_b\}$ is its decomposition then, for each $\lambda \in [0, 1]$, S_λ is an isolated invariant set with respect to f_λ , $\{S_{b\lambda}\} = \{S_{b\lambda}\}_{b \in B}$ is its decomposition and $h(\{S_{b\lambda}\}, f_\lambda, X)$ does not depend on λ .*

Proof. It is enough to show that each $\lambda \in [0, 1]$ admits a neighborhood U such that $h(\{S_{b\mu}\}, f_\mu, X)$ is constant in $\mu \in U$. Let N be an isolating neighborhood for S_λ with respect to f_λ which admits a decomposition $\{N_b\}$ such that $N_b \cap S_\lambda = S_{b\lambda}$ for each $b \in B$. As a consequence of the existence theorem for index pairs for multivalued upper semicontinuous maps (see [6], theorem 2.6) we obtain the existence of a stable index pair, i. e. a pair $Q = (Q_1, Q_0)$ which is an index pair for S_μ with respect to f_μ for each μ in a certain interval J being a neighborhood of λ in $[0, 1]$, such that $Q_1 \subset N$. Let $D_b = cl(Q_1 \setminus Q_0) \cap N_b$ for each $b \in B$. Since $D_b \cap S_\lambda = S_{b\lambda}$, by making J smaller if necessary we may assume that

$$D_b \cap S_\mu = S_{b\mu}$$

for each $\mu \in J$. This means that Q is compatible with the decomposition $\{S_{b\mu}\}$ of S_μ . Furthermore, since J is an interval, the homotopy class of the index map $(f_\mu)_Q : Q_1/Q_0 \rightarrow Q_1/Q_0$ does not depend on $\mu \in J$. By definition 3.3, the index object $I(Q, \{D_b\}, f_\mu)$ is independent of $\mu \in J$. \square

The next theorem shows that the Conley index for a decomposition $\{S_b\}$ carries information about ordinary Conley indices of some subsets of S .

Theorem 4.2 *Let $\{S_b\}$ be a decomposition of an isolated invariant set S with respect to a continuous map $f : X \rightarrow X$. For each sequence $\bar{Y} = (Y_0, Y_1, \dots, Y_{n-1})$ of members of A we put $\tilde{S}_{\bar{Y}} = \bigcap_{i=0}^{n-1} f^{-i}(S_{Y_i})$ and $S_{\bar{Y}} = \text{Inv}_{f^n}(\tilde{S}_{\bar{Y}})$. $S_{\bar{Y}}$ is an isolated invariant set with respect to f^n contained in S and*

$$h(S_{\bar{Y}}, f^n, X) = \mathcal{P}_{i(\bar{Y})}(h(\{S_b\}, f, X))$$

Proof. Let N be an isolating neighborhood for S with respect to f admitting a decomposition $\{N_b\}$ such that $N_b \cap S = S_b$ for each $b \in B$. By proposition 2.1 in [20], $N_{\bar{Y}} = \bigcap_{i=0}^{n-1} f^{-i}(N_{Y_i})$ is an isolating neighborhood with respect to f^n and its invariant part is contained in S . Therefore, $\text{Inv}_{f^n}(N_{\bar{Y}}) = \text{Inv}_{f^n}(\tilde{S}_{\bar{Y}}) = S_{\bar{Y}}$, which means that $N_{\bar{Y}}$ is an isolating neighborhood for $S_{\bar{Y}}$ with respect to f^n . We have completed the proof of the first part of the theorem. In order to see that the formula for the Conley index of $S_{\bar{Y}}$ holds we shall make use of the following fact (see [20], lemma 3.1).

If $Q = (Q_1, Q_0)$ is a regular index pair for S such that $Q_1 \subset N$ then

$$h(S_{\bar{Y}}, f^n, X) = [Q_1/Q_0, f_{(Q, \{D_b\})}^{Y_{n-1}} \circ f_{(Q, \{D_b\})}^{Y_{n-2}} \circ \dots \circ f_{(Q, \{D_b\})}^{Y_0}]$$

where $D_b = \text{cl}(Q_1 \setminus Q_0) \cap N_b$ for each $b \in B$.

Since such index pairs exist, this formula together with the definition of the \mathcal{P} -type functors prove the theorem. \square

Theorem 4.3 (Locality) *If $f : X \rightarrow X$ and $g : X \rightarrow X$ are continuous maps, S is an isolated invariant set with respect to f and f and g are equal on a certain neighborhood of S then S is an isolated invariant set with respect to g and for any its decomposition $\{S_b\}$,*

$$h(\{S_b\}, f, X) = h(\{S_b\}, g, X).$$

Proof. By proposition 3.1, there exists an index pair $Q = (Q_1, Q_0)$ for S compatible with the decomposition $\{S_b\}$ such that f and g restricted to Q_1 are equal. Since the corresponding index objects depend only on the restrictions of both maps to Q_1 , they are the same. \square

The rest of this section is devoted to the formulation and proof of the Wazewski property of the Conley index for decompositions of isolated invariant sets and a bound for the number of periodic points of f in terms of the Conley index for decompositions. Fix a locally compact metric space X , a continuous map f of X into itself, an isolated invariant set S and its decomposition $\{S_b\}$. Let N be a fixed isolating neighborhood for S admitting a decomposition $\{N_b\}$ such that $N_b \cap S = S_b$. Let

$$S^+ = \text{Inv}^+ N = \{x \in N : \forall_{i \in \mathbb{Z}^+} f^i(x) \in N\}.$$

The map $p : S^+ \rightarrow \Pi = \prod_{i \in \mathbb{Z}^+} B$ is defined by

$$p(x) = (\eta(f^i(x)))_{i=0}^{\infty} \tag{4.1}$$

where $\eta : S^+ \rightarrow B$ is defined by

$$\eta(x) = b \quad \text{if and only if} \quad x \in N_b$$

Clearly, both p and η are continuous if we topologize B with the discrete topology. Furthermore, $p \circ f = \sigma \circ p$ where $\sigma : \Pi \rightarrow \Pi$ is the shift map. This means that p is a semiconjugacy onto its image. In what follows, we shall give lower bounds for the image of p and the image of the set of periodic points of f under p in terms of the Conley index for decompositions.

Definition 4.1 *Let $(Y, \{g^Z\}_{Z \in A})$ be an object in a category $\mathcal{K}_{[A]}$ ($\mathcal{K} = \mathcal{Htop}, \mathcal{M}$ or \mathcal{M}_G). We put*

$$\Pi_0(Y, \{g^Z\}_{Z \in A}) = \{(b_i)_{i=0}^{\infty} \in \Pi : \forall_{n \in \mathbb{Z}^+} g^{\{b_n\}} \circ g^{\{b_{n-1}\}} \circ \dots \circ g^{\{b_0\}} \neq 0\},$$

$$\begin{aligned}\Pi(Y, \{g^Z\}_{Z \in A}) &= \bigcap_{n \in \mathbb{Z}^+} \sigma^n(\Pi_0(Y, \{g^Z\}_{Z \in A})), \\ \Pi_0^*(Y, \{g^Z\}_{Z \in A}) &= \{(b_i)_{i=0}^\infty \in \Pi : \forall n \in \mathbb{Z}^+ \quad g^{\{b_0\}} \circ g^{\{b_1\}} \circ \dots \circ g^{\{b_n\}} \neq 0\}, \\ \Pi^*(Y, \{g^Z\}_{Z \in A}) &= \bigcap_{n \in \mathbb{Z}^+} \sigma^n(\Pi_0^*(Y, \{g^Z\}_{Z \in A})).\end{aligned}$$

The next proposition collects some of the properties of the sets defined above.

Proposition 4.1 *Let $(Y, \{g^Z\}_{Z \in A})$ and $(\bar{Y}, \{\bar{g}^Z\}_{Z \in A})$ be objects in $\mathcal{K}_{[A]}$.*

- 1°. $\Pi_0(Y, \{g^Z\}_{Z \in A})$ and $\Pi_0^*(Y, \{g^Z\}_{Z \in A})$ are compact,
- 2°. $\sigma(\Pi_0(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0(Y, \{g^Z\}_{Z \in A})$
and $\sigma(\Pi_0^*(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0^*(Y, \{g^Z\}_{Z \in A})$,
- 3°. if $(Y, \{g^Z\}_{Z \in A})$ and $(\bar{Y}, \{\bar{g}^Z\}_{Z \in A})$ are isomorphic in $\mathcal{K}_{[A]}$ then

$$\exists_{n \in \mathbb{Z}^+} \quad \sigma^n(\Pi_0(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0(\bar{Y}, \{\bar{g}^Z\}_{Z \in A})$$

and

$$\exists_{m \in \mathbb{Z}^+} \quad \sigma^m(\Pi_0^*(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0^*(\bar{Y}, \{\bar{g}^Z\}_{Z \in A}).$$

Therefore,

$$\Pi(Y, \{g^Z\}_{Z \in A}) = \Pi(\bar{Y}, \{\bar{g}^Z\}_{Z \in A})$$

and

$$\Pi^*(Y, \{g^Z\}_{Z \in A}) = \Pi^*(\bar{Y}, \{\bar{g}^Z\}_{Z \in A}).$$

In particular, the sets $\Pi(I)$ and $\Pi^*(I)$ can be defined in the obvious way for isomorphism classes I in $\mathcal{K}_{[A]}$.

- 4°. Suppose that \mathcal{K} and \mathcal{L} are categories, each of them equal to \mathcal{Htop} , \mathcal{M} or \mathcal{M}_G and $F : \mathcal{K} \rightarrow \mathcal{L}$ is a functor mapping the zero morphisms into the zero morphisms. If F is covariant then

$$\Pi_0(F_{[A]}(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0(Y, \{g^Z\}_{Z \in A})$$

and

$$\Pi_0^*(F_{[A]}(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0^*(Y, \{g^Z\}_{Z \in A}).$$

If F is contravariant then

$$\Pi_0^*(F_{[A]}(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0(Y, \{g^Z\}_{Z \in A})$$

and

$$\Pi_0(F_{[A]}(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0^*(Y, \{g^Z\}_{Z \in A}).$$

The same inclusions hold for Π_0 replaced with Π .

Proof. We shall prove only the parts concerning $\Pi(Y, \{g^Z\}_{Z \in A})$ and $\Pi_0(Y, \{g^Z\}_{Z \in A})$. The proofs of the other parts of the proposition are dual.

1°. If $(b_i)_{i=0}^\infty \notin \Pi_0(Y, \{g^Z\}_{Z \in A})$ then, for some $n \in Z^+$,

$$g^{\{b_n\}} \circ g^{\{b_{n-1}\}} \circ \dots \circ g^{\{b_0\}} = 0$$

Hence

$$\prod_{i=0}^n \{b_i\} \times \prod_{i=n+1}^\infty B \subset \Pi \setminus \Pi_0(Y, \{g^Z\}_{Z \in A}).$$

Thus, the complement of $\Pi_0(Y, \{g^Z\}_{Z \in A})$ is open in Π so that this set is compact.

2°. If $(b_i)_{i=0}^\infty \in \Pi_0(Y, \{g^Z\}_{Z \in A})$ then, for each $n \in Z^+$,

$$g^{\{b_{n+1}\}} \circ g^{\{b_n\}} \circ \dots \circ g^{\{b_0\}} \neq 0.$$

Hence,

$$g^{\{b_{n+1}\}} \circ g^{\{b_n\}} \circ \dots \circ g^{\{b_1\}} \neq 0.$$

It follows that

$$(b_{i+1})_{i=0}^\infty = \sigma((b_i)_{i=0}^\infty) \in \Pi_0(Y, \{g^Z\}_{Z \in A}).$$

3°. Let

$$[\{h^{\bar{Z}}\}_{\bar{Z} \in A^n}, n] : (Y, \{g^Z\}_{Z \in A}) \longrightarrow (\bar{Y}, \{\bar{g}^Z\}_{Z \in A})$$

and

$$[\{l^{\bar{Z}}\}_{\bar{Z} \in A^m}, m] : (\bar{Y}, \{\bar{g}^Z\}_{Z \in A}) \longrightarrow (Y, \{g^Z\}_{Z \in A})$$

be reciprocal isomorphisms. Then, by the definition of the $\mathcal{K}_{[A]}$ category and definition 4.1, for each $(b_i)_{i=0}^\infty \in \Pi_0(Y, \{g^Z\}_{Z \in A})$ and sufficiently large $k \in Z^+$,

$$\begin{aligned} 0 &\neq g^{\{b_{n+m+k-1}\}} \circ \dots \circ g^{\{b_0\}} = \\ &= l^{\{b_{n+m+k-1}, \dots, b_{n+k}\}} \circ h^{\{b_{n+k-1}, \dots, b_k\}} \circ g^{\{b_{k-1}\}} \circ \dots \circ g^{\{b_0\}} = \\ &= l^{\{b_{n+m+k-1}, \dots, b_{n+k}\}} \circ \bar{g}^{\{b_{n+k-1}\}} \circ \dots \circ \bar{g}^{\{b_n\}} \circ h^{\{b_{n-1}, \dots, b_0\}}. \end{aligned}$$

Hence, $g^{\{b_{n+k-1}\}} \circ \dots \circ g^{\{b_n\}} \neq 0$ and therefore $\sigma^n((b_i)_{i=0}^\infty) \in \Pi_0(\bar{Y}, \{\bar{g}^Z\}_{Z \in A})$.

The second part of 3° follows immediately from the first one, proposition 4.1, 2° and definition 4.1.

4° follows immediately from definition 4.1. \square

Theorem 4.4 (Ważewski property) *Let $(Y, \{g^Z\}_{Z \in A})$ be an object in $\mathcal{Htop}_{[A]}$ such that*

$$h(\{S_b\}, f, X) = [Y, \{g^Z\}_{Z \in A}].$$

Then:

1°. *There exists $n \in Z^+$ such that*

$$\sigma^n(\Pi_0(Y, \{g^Z\}_{Z \in A})) \subset p(S^+),$$

2°. $\Pi(Y, \{g^Z\}_{Z \in A}) \subset p(S)$.

Proof. Let $Q = (Q_1, Q_0)$ be an index pair for S such that $Q_1 \subset N$. Let $D_b = cl(Q_1 \setminus Q_0) \cap N_b$. Clearly, $D_b \cap S = S_b$. The index object $I(Q, \{D_b\})$ is isomorphic to $(Y, \{g^Z\}_{Z \in A})$. Hence, by proposition 4.1, 3°,

$$\exists_{n \in Z^+} \sigma^n(\Pi_0(Y, \{g^Z\}_{Z \in A})) \subset \Pi_0(I(Q, \{D_b\})).$$

It follows that to prove 1° it is enough to show that the set on the right-hand side is contained in $p(S^+)$. Take $(b_i)_{i=0}^\infty \in \Pi_0(I(Q, \{D_b\}))$. Then, for each

$k \in Z^+$ there exists an $x_k \in Q_1$ such that

$$f_{(Q, \{D_b\})}^{\{b_k\}} \circ \dots \circ f_{(Q, \{D_b\})}^{\{b_0\}}([x_k]) \neq [Q_0].$$

The formula (3.1) proves that

$$\forall_{i \in \{0, \dots, k\}} f^i(x_k) \in D_{b_i} \cap (Q_1 \setminus Q_0).$$

Now, let x_* be an accumulation point of the sequence $\{x_k\}$. Clearly, $f^i(x_*) \in D_{b_i} \subset N$ for all $i \in Z^+$. Thus, $x_* \in S^+$ and $p(x_*) = (b_i)_{i=0}^\infty$. The proof of 1° is complete. To prove 2°, notice that

$$\sigma^{k+n}(\Pi_0(Y, \{g^Z\}_{Z \in A})) \subset \sigma^k(p(S^+)) = p(f^k(S^+))$$

for each $k \in Z^+$. Hence,

$$\Pi(Y, \{g^Z\}_{Z \in A}) = \bigcap_{k \in Z^+} \sigma^{k+n}(\Pi_0(Y, \{g^Z\}_{Z \in A})) \subset \bigcap_{k \in Z^+} p(f^k(S^+)).$$

Since $\{f^k(S^+)\}$ is a decreasing sequence of compact sets intersecting at S , the set on the right-hand side is equal to $p(S)$. \square

Theorem 4.5 (Detection of periodic points) *Let $(b_i)_{i=0}^\infty \in \Pi$ be T -periodic. Put $\bar{Y} = (\{b_0\}, \{b_1\}, \dots, \{b_{T-1}\}) \in A^T$. If, for some positive integer n ,*

$$\Lambda(\mathcal{P}_{\bar{Y}^n}(h^*(\{S_b\}, f, X))) \neq 0 \tag{4.2}$$

then there exists an nT -periodic point $x \in S$ of f such that $p(x) = (b_i)_{i=0}^\infty$. If (4.2) holds for $n = 1$ and T is the principal period of $(b_i)_{i=0}^\infty$ then x can be chosen in such a way that T is its principal period.

Proof. By (4.2), theorem 4.2 and remark 2.1,

$$\Lambda(h^*(S_{\bar{Y}^n}, f^{nT}, X)) \neq 0.$$

By theorem 1.1, there exists an $x \in S_{\bar{Y}^n}$ such that $f^{nT}(x) = x$. Since $p(S_{\bar{Y}^n}) = \{(b_i)_{i=0}^\infty\}$, this proves the first part of the theorem. Since $p \circ f = \sigma \circ p$, the principal period of $p(x)$ must not exceed the principal period of x with respect to f . Thus, if $n = 1$ then the principal period of x equals T . \square .

5. Horseshoes

In many chaotic dynamical systems arising from differential equations a kind of behaviour resembling that of the Smale's horseshoe map is observed (see [8], [22]). In this section we provide an example of a criterion for chaos based on the index defined in section 3 and compute the indices of decompositions of invariant sets of the Smale's horseshoe (U-horseshoe) and the G-horseshoe maps. In the sequel, we deal only with decompositions of isolated invariant sets into two disjoint subsets, i.e. we set $B = \{1, 2\}$. By

$$\begin{bmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ \dots & \dots & \dots & \dots \\ a_{k,1} & a_{k,2} & \dots & a_{k,n} \\ \hline a_{k+1,1} & a_{k+1,2} & \dots & a_{k+1,n} \\ \dots & \dots & \dots & \dots \\ a_{n,1} & a_{n,2} & \dots & a_{n,n} \end{bmatrix}$$

we denote the class of all objects in $\mathcal{M}_{[A]}$ isomorphic to the object

$(\Xi^n, \{\psi^Z\}_{Z \in A})$ such that the matrices of $\psi^{\{1\}}$ and $\psi^{\{2\}}$ in the standard basis are

$$\begin{bmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ \dots & \dots & \dots & \dots \\ a_{k,1} & a_{k,2} & \dots & a_{k,n} \\ 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 \\ a_{k+1,1} & a_{k+1,2} & \dots & a_{k+1,n} \\ \dots & \dots & \dots & \dots \\ a_{n,1} & a_{n,2} & \dots & a_{n,n} \end{bmatrix}$$

(respectively) and $\psi^Z = \sum_{b \in Z} \psi^{\{b\}}$ for each $Z \in A$. Let us note the following

Corollary 5.1 *Let f be a continuous map of a locally compact metric space X into itself, S an isolated invariant set with respect to f and $\{S_b\}$ its de-*

composition. Suppose that, for some $q \in Z^+$,

$$h^q(\{S_b\}, f, X) = \begin{bmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{bmatrix}.$$

where $a_{i,j} \in \Xi$ are nonzero. If Ξ is a domain then the map $p : S \rightarrow \Pi$ defined by (4.1) is a surjection. If, in addition,

$$h^r(\{S_b\}, f, X) = 0$$

for each $r \neq q$, Ξ is the field of rational numbers and X is an ENR then each periodic sequence in Π is an image (under p) of a periodic point of f in S of the same principal period.

Proof. Let $\psi_1, \psi_2 : \Xi^2 \rightarrow \Xi^2$ have matrices $\begin{bmatrix} a_{1,1} & a_{1,2} \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & 0 \\ a_{2,1} & a_{2,2} \end{bmatrix}$ in the standard basis. For each sequence $(b_0, b_1, \dots, b_{T-1})$ of members of B ,

$$\psi_{b_{T-1}} \circ \psi_{b_{T-2}} \circ \dots \circ \psi_{b_0} \neq 0.$$

More precisely, the matrix of the composition on the left-hand side has one row of zeros and one row of nonzero members of Ξ (in particular, its trace is nonzero if Ξ is a field: we shall use this fact later). Thus,

$$\Pi^* \left(\begin{bmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{bmatrix} \right) = \Pi.$$

Applying proposition 4.1,4° for F being the q -dimensional cohomology functor,

$$\Pi(h(\{S_b\}, f, X)) = \Pi.$$

To finish the proof of the first part, apply theorem 4.4,2°. In order to prove the remaining part, take a sequence $(b_i)_{i=0}^\infty \in \Pi$ of principal period T . Let $\bar{Y} = (\{b_0\}, \{b_1\}, \dots, \{b_{T-1}\})$. By assumptions, $tr(\mathcal{P}_{\bar{Y}}(h^r(\{S_b\}, f, X))) = 0$ for all $r \neq q$. Since the trace is nonzero for $r = q$, the Lefschetz number of

$\mathcal{P}_{\bar{Y}}(h^*(\{S_b\}, f, X))$ is nonzero. Theorem 4.5 implies that $(b_i)_{i=0}^\infty$ is an image of a periodic point of f of principal period T . \square

In a moment we shall show that the assumptions of the above corollary are satisfied by the G-horseshoe and the U-horseshoe maps. We note that, by the continuation property of the Conley index for decompositions, corollary 6.1 generalizes theorem 2.4 in [9] and theorem 2.1 in [23].

Example (cf [11]). Let $G : R^2 \rightarrow R^2$ be a G-horseshoe map, i.e. a map of R^2 into itself being affine on each of the rectangles $A_i B_i C_i D_i$ ($i=1,2$). The rectangles and their images under G are depicted on figure 1.




Fig.1. G-horseshoe

We are interested in the Conley index of the decomposition $\{S_b\}$ of the maximal invariant set S in $A_1B_1C_1D_1 \cup A_2B_2C_2D_2$, where $S_b = S \cap A_iB_iC_iD_i$ for each $b \in B = \{1, 2\}$. By theorem 4.3 we can assume that G maps the rectangle $A_2B_2C_1D_1$ outside the rectangle $A_1B_1C_2D_2$. Let Q_0 be the set of all points in the rectangle $A_1B_1C_2D_2$ mapped by G outside its interior (it is the shadowed region being a disjoint union of three rectangles - see fig.1). One can easily see that if we put $Q_1 = A_1B_1C_2D_2$ then the pair $Q = (Q_1, Q_0)$ is an index pair for S , compatible with the decomposition $\{S_b\}$ of S . Clearly, the quotient space Q_1/Q_0 has the homotopy type of the wedge sum of two pointed circles. Thus,

$$H^q(Q_1/Q_0) = \begin{cases} \Xi^2 & \text{if } q = 1 \\ 0 & \text{otherwise} \end{cases} .$$

In $H^1(Q_1/Q_0)$ we can chose the basis consisting of vectors e_b ($b \in B$), being the images of the generators of $H^1(A_1B_1C_2D_2/(A_{1-b}B_{1-b}C_{1-b}D_{1-b} \cup Q_0))$ under the inclusion-induced homomorphism. Then, $H^1(G_Q)$ maps e_b into $e_1 + e_2$ or $e_1 + e_2 - 2e_{1-b}$ according to the choice of generators. Hence (notice that $H^1(r^Z)$ is the natural projection onto the submodule generated by $\{e_b : b \in Z\}$),

$$h^1(\{S_b\}, G, R^2) = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} .$$

Clearly, the cohomological indices in other dimensions are trivial. In the same way, one can prove that if U is the U-horseshoe map (cf [11]) and $\{S_b\}$ is the decomposition of its invariant set obtained in the analogous way then

$$h^1(\{S_b\}, U, R^2) = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$$

and the cohomological indices of all other dimensions are trivial.

References

- [1] Ch.Conley, *Isolated Invariant Sets and the Morse Index*, CBMS Regional Conference Series in Mathematics, Vol. 38, AMS Publications, Providence 1978.
- [2] A.Dold, *Introduction to Algebraic Topology*, Springer-Verlag 1972.
- [3] R.Franzosa, *Index filtrations and the homology index braid for partially ordered Morse decompositions*, Trans. Am. Math. Soc. 298(1986),193-213.
- [4] R.Franzosa, *The connection matrix theory for Morse decompositions*, Trans. Am. Math. Soc. 311(1989),561-592.
- [5] R.Franzosa, *The continuation theory for Morse decompositions and connection matrices* Trans. Am. Math. Soc. 310(1988),781-803.
- [6] T.Kaczyński and M.Mrozek, *Conley index for discrete multivalued dynamical systems* , to appear in Topology Appl.
- [7] Ch.McCord, K.Mischaikow and M.Mrozek, *Zeta functions, periodic trajectories and the Conley index*, preprint CDSNS93-152.
- [8] K.Mischaikow and M.Mrozek, *Chaos in Lorenz Equations: a Computer Assisted Proof*, preprint CDSNS93-124.
- [9] K.Mischaikow and M.Mrozek, *Isolating neighborhoods and chaos*, preprint CDSNS93-116.
- [10] M.Mrozek, *Index pairs and the fixed point index for semidynamical systems with discrete time*, Fund. Math. 133(1989),179-194.
- [11] M.Mrozek, *Leray functor and the cohomological Conley index for discrete dynamical systems*, Trans. Am. Math. Soc. 318(1990),149-178.
- [12] M.Mrozek, *Open index pairs, the fixed point index and rationality of zeta functions*, Ergodic Th.Dyn.Sys 10(1990),555-564.
- [13] M.Mrozek, *Shape index and other indices of Conley type for local maps on locally compact Hausdorff spaces*, preprint CDSNS92-106.
- [14] M.Mrozek, *The Conley index on compact ANR's is of finite type*, Results in Mathematics 18(1990),306-313.
- [15] M.Mrozek, *The Morse equation in the Conley's index theory for homeomorphisms*, Topology Appl. 38(1991),45-60.
- [16] M.Mrozek and K.P.Rybakowski, *A cohomological Conley index for maps on metric spaces*, J.Diff.Equations 90(1991),143-171.
- [17] J.W.Robbin and D.Salamon, *Dynamical systems, shape theory and the Conley index*, Ergodic Th. Dyn. Sys. 8 Suppl(1988),375-393.
- [18] D.Salamon, *Connected simple systems and the Conley index of isolated invariant sets*, Trans. Am. Math. Soc. 291(1985),1-41.
- [19] R.Srzednicki, *Fixed point index of iterations and detection of chaos*, preliminary report.
- [20] A.Szymczak, *The Conley index and symbolic dynamics*, to appear in Topology.
- [21] A.Szymczak, *The Conley index for discrete semidynamical systems*, to appear in Topology Appl.
- [22] S.Wiggins, *Global Bifurcations and Chaos*, Springer-Verlag 1988.
- [23] P.Zgliczyński *Fixed point index for iterations of maps, topological horseshoe and chaos*, preprint.

Institute of Mathematics,
Polish Academy of Sciences,
ul. Abrahama 18,
81-825 Sopot,
Poland.
E-mail: matacs@halina.univ.gda.pl