


PIC Math: A Modified SEIR Model for the Spread of Ebola in Western Africa

Paul Diaz, Eric Jones, Kelsey Kalmbach, and Chelsea Sandridge

Colorado School of Mines

April 28, 2015

Background - Cumulative Ebola Outbreak Distribution Map




Goal: Mitigate the spread of Ebola in West Africa by determining the optimal placement of a treatment center

Model Assumptions

- ▶ Infected individuals can move to three different compartments: Removed and infectious, removed and buried, removed and recovered
- ▶ Individuals who have died from the disease but who have not yet been buried can still infect susceptible individuals
- ▶ Individuals who recover from the disease are no longer susceptible
- ▶ Once hospitalized, infected individuals can still infect others. However, those who die in the hospital are buried immediately, and thus cannot infect others once dead
- ▶ Hospitals have unlimited space, but there is some delay in hospitalization

State Diagram



The Model

$$\left\{ \begin{array}{l} \frac{dS}{dt} = \alpha S - \beta_1 SI - \beta_2 SR_I - \beta_3 SH \\ \frac{dE}{dt} = \beta_1 SI + \beta_2 SR_I + \beta_3 SH - \delta E \\ \frac{dI}{dt} = \delta E - \gamma_1 I - \psi I \\ \frac{dH}{dt} = \psi I - \gamma_2 H \\ \frac{dR_I}{dt} = \rho_1 \gamma_1 I - \omega R_I \\ \frac{dR_B}{dt} = \omega R_I + \rho_2 \gamma_2 H \\ \frac{dR_R}{dt} = (1 - \rho_1) \gamma_1 I + (1 - \rho_2) \gamma_2 H \end{array} \right.$$

Parameter Fitting


Data: country time series data compiled from WHO reports up to December, 2014 [9].

fminsearch: Finds the minimum of unconstrained multivariable function using derivative-free method.


Error function: Sums of Squares

$$e := \sum \sqrt{(R_{data} - (R_I + R_B))^2 + (I_{data} - (I + H))^2}$$


Parameter Fitting




Parameter Fitting



Parameter Fitting




Parameter Fitting




Adding Transportation Between Countries

$$\left\{ \begin{array}{l} \frac{dS_i}{dt} = \alpha S_i - \beta_1 S_i I_i - \beta_2 S_i R_{I,i} - \beta_3 S_i H_i \\ \frac{dE_i}{dt} = \beta_1 S_i I_i + \beta_2 S_i R_{I,i} + \beta_3 S_i H_i - \delta E_i - \phi_{ij} E_i + \phi_{ji} E_j \\ \frac{dI_i}{dt} = \delta E_i - \gamma_1 I_i - \psi I_i \\ \frac{dH_i}{dt} = \psi I_i - \gamma_2 H_i \\ \frac{dR_{I,i}}{dt} = \rho_1 \gamma_1 I_i - \omega R_{I,i} \\ \frac{dR_{B,i}}{dt} = \omega R_{I,i} + \rho_2 \gamma_2 H_i \\ \frac{dR_{R,i}}{dt} = (1 - \rho_1) \gamma_1 I_i + (1 - \rho_2) \gamma_2 H_i \end{array} \right.$$


Coupled Parameter Fitting



Coupled Parameter Fitting



Coupled Parameter Fitting



Coupling Method and Results

- ▶ Created coupled (21 compartment) model, with initial conditions “stitched” together
- ▶ Used previously fit parameter values for each country, then fit transportation coefficients to the same data as before
- ▶ **Error without coupling: 142**
- ▶ **Error with coupling: 14**

Placement Based on Parameter Values


	Guinea	Liberia	Sierra Leone
(hospitalization rate) ⁻¹ ψ	1.44	0.497	2.21
death rate (not hospitalized) ρ_1	0.649	0.77	0.451
death rate (hospitalized) ρ_2	0.59	0.7	0.41

Placement Based on Sensitivity Analysis

Relative change in population steady state given a change in a parameter value

$\frac{\partial I}{\partial \psi} \Delta \psi$	
Guinea	-336.882
Liberia	-84.6367
Sierra Leone	-0.467521

Placement Based on Minimizing Infectious Population



Future Work

- ▶ Increase resolution — provinces rather than countries
- ▶ Assume hospitals have a finite capacity (logistic transfer rate)
- ▶ Observe epidemic by placing “patient zero” in one country and watching it spread with the coupled model
- ▶ Gather recent data and update our model while the epidemic is ongoing

Questions?

Variable Definitions

Populations:

S = Susceptible Population

E = Exposed Population (infected but not yet showing symptoms)

I = Infectious Population

H = Hospitalized Population

R_I = Removed and Infectious Population (dead but not yet buried)

R_B = Removed and Buried

R_R = Removed and Recovered

Variable Definitions

Parameters:

α = population growth constant [3]

β_1 = transmission rate between infected and susceptible

β_2 = transmission rate between removed and still infectious and susceptible

β_3 = transmission rate between hospitalized and susceptible

δ = rate at which people move from exposed to infected [4]

γ_1 = (average time with disease for unhospitalized individuals) $^{-1}$

γ_2 = (average time with disease for hospitalized individuals) $^{-1}$

ψ = (average time that people become hospitalized) $^{-1}$

$\rho_1 = 1.1 \times \rho_2$ = the proportion of people who die of the disease who are not hospitalized [5]

ρ_2 = the proportion of people who die of the disease who are hospitalized [5]

ω = (time until one is buried) $^{-1}$

References

- [1] Center for Disease Control. *Questions and Answers: Estimating the Future Number of Cases in the Ebola Epidemic*, 2015.
- [2] J. Lewnard et. al. *Dynamics and Control of Ebola virus Transmission in Montserrado, Liberia: a mathematical modeling analysis* Department of Epidemiology of Microbial Diseases. Yale, CT. Dec 2014. <http://www.cdc.gov/vhf/ebola/outbreaks/2014-west-africa/qa-mmwr-estimating-future-cases.html>
- [3] "Population Growth (annual %)." The World Bank. Web. 26 Mar. 2015. <http://data.worldbank.org/indicator/SP.POP.GROW>.
- [4] "Signs and Symptoms." Centers for Disease Control and Prevention. N.p., n.d. Web. 27 Mar. 2015. <http://www.cdc.gov/vhf/ebola/symptoms/index.html>.
- [5] "Case Fatality Rate for Ebolavirus." *Epidemic : Molecular Epidemiology and Evolution of Viral Pathogens*. Web. 17 Apr. 2015. <http://epidemic.bio.ed.ac.uk/ebolavirus-fatality-rate>.
- [6] Nielsen, Carrie F., Sarah Kidd, Ansumana Sillah, Edward Davis, and Jonathan Mermin. "Improving Burial Practices and Cemetery Management During an Ebola Virus Disease Epidemic — Sierra Leone, 2014." Centers for Disease Control and Prevention. *Centers for Disease Control and Prevention*, 16 Jan. 2015. Web. 26 Mar. 2015.
- [7] Epatko, Larisa. "70 percent Ebola death rate? Here's how they calculate it." *PBS News Hours*. N.p., 16 Oct. 2014. Web. 26 Mar. 2015. <http://www.pbs.org/newshour/rundown/70-percent-ebola-death-rate-calculate/>.
- [8] Shaby, Ben. "Fitting the SIR model to data" *Statistical and Applied Mathematical Sciences Institute Undergraduate Workshop*. 18 May, 2010. http://www.samsi.info/sites/default/files/Shaby_sir_lab.pdf
- [9] Rivers, Caitlin. "Consolidated, machine-readable ebola situation reports" *Virginia Tech Network Dynamics and Simulation Science Laboratory*. 31 Dec, 2014. <https://github.com/muxspace/ebola.sitreps>