

- The weak interactions have many peculiar features that set them apart from QCD and EM:
- i) Every single matter particle in the SM exhibits weak interactions (only charged for EM, only quarks for QCD)
 - ii) The force mediators are massive (unlike photons and gluons)
 - iii) The weak interactions violate parity, charge conjugation and CP.
 - iv) The weak interactions can change flavor, i.e. particle type \Rightarrow responsible for decays!

Perhaps the strangest part of the weak interactions is that they are not realized as a symmetry of the SM, at least not at room temperature type energies. We will eventually explain what this means and in fact this will solve the mediator mass issue by bringing in the Higgs mechanism. But more on that later.

To keep in line with our development of EM and QCD as theories of local (or gauge) invariance, we will go ahead and formulate the weak interactions in terms of a gauge symmetry. This is relevant since at some point in the history of the universe this is how it appeared. More on that later!

Perhaps the most surprising feature of formulating the weak interactions in terms of a gauge symmetry is that to do so, we are forced to "unify" the weak force with EM!

Electroweak Unification

It is often said that the total gauge symmetry group of the SM is $\underbrace{SU(3)}_{\text{QCD}} \times \underbrace{SU(2)}_{\text{Weak}} \times \underbrace{U(1)}_{\text{E+H}}$

However this is not quite right. The correct groups are:

$$\begin{array}{ccc} \underbrace{SU(3)}_{\text{QCD}} & \underbrace{SU(2)_L \times U(1)_Y}_{\text{Electroweak}} & \text{High Energy} \\ & \Downarrow \begin{array}{l} \text{left} \\ \text{chiral} \end{array} & \uparrow \text{hypercharge} \\ \underbrace{SU(3)}_{\text{QCD}} & \underbrace{U(1)_{EM}}_{\text{E+H}} & \text{Low Energy} \end{array}$$

So we need to start with $SU(2)_L \times U(1)_Y$. The "L" in $SU(2)_L$ means that this symmetry is only relevant for "left-handed" fermion states. This is a bit of a misnomer since handedness has to do with helicity, whereas in actuality the $SU(2)$ acts on states of definite chirality (which does match helicity for massless particles).

Recall: $\mathcal{L}_{\text{Dirac}} = (k_c) \bar{\psi} \gamma^\mu \partial_\mu \psi + m c^2 \bar{\psi} \psi$

$$= -k_c \left(\psi_-^\dagger \partial_\mu \bar{\psi}^\mu \psi_- + \psi_+^\dagger \partial_\mu \bar{\psi}^\mu \psi_+ \right) + m c^2 (\psi_-^\dagger \psi_+ + \psi_+^\dagger \psi_-) \quad \left. \vphantom{\begin{matrix} \psi_-^\dagger \\ \psi_+^\dagger \end{matrix}} \right\} \begin{matrix} \text{This is in terms} \\ \text{of 2-component objects} \\ \psi_+, \psi_- \text{ and c.c.} \end{matrix}$$

$\uparrow \qquad \qquad \qquad \uparrow$
 $\bar{\psi}^\mu = (I, \vec{\sigma}^i) \qquad \bar{\psi}^\mu = (I, -\vec{\sigma}^i)$

To work w/ 4 component objects consider $\psi_R = \begin{pmatrix} \psi_+ \\ 0 \end{pmatrix}$ and $\psi_L = \begin{pmatrix} 0 \\ \psi_- \end{pmatrix}$. Then $\psi = \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix} = \psi_R + \psi_L$

But recall that $P_\pm = \frac{1}{2}(1 \pm \gamma^5) \Rightarrow P_+ \psi = \frac{1}{2} \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \psi_R \\ \psi_L \end{pmatrix} = \psi_R, P_- \psi = \psi_L$

However for $\bar{\psi} = i \psi^\dagger \gamma^0 \Rightarrow \bar{\psi}_R = \overline{\left(\frac{1+\gamma^5}{2} \psi \right)} = i \left(\frac{1+\gamma^5}{2} \psi \right)^\dagger \gamma^0 = i \psi^\dagger \frac{1+\gamma^5}{2} \gamma^0 = i \psi^\dagger \gamma^0 \frac{1-\gamma^5}{2} = \bar{\psi} \frac{1-\gamma^5}{2}$

Then we can write: $\mathcal{L}_{\text{Dirac}} = (k_c) (\bar{\psi}_R + \bar{\psi}_L) \gamma^\mu \partial_\mu (\psi_R + \psi_L) + m c^2 (\bar{\psi}_R + \bar{\psi}_L) (\psi_R + \psi_L)$

Note that: $\bar{\psi}_R \gamma^\mu \psi_L = \bar{\psi} \frac{1-\gamma^5}{2} \gamma^\mu \frac{1-\gamma^5}{2} \psi = \bar{\psi} \gamma^\mu \frac{1+\gamma^5}{2} \frac{1-\gamma^5}{2} \psi = \bar{\psi} \gamma^\mu \frac{1-\gamma^5}{2} \psi = 0$
 \Rightarrow All derivative terms mixing L w/ R vanish.

$\bar{\psi}_R \psi_R = \bar{\psi} \frac{1-\gamma^5}{2} \frac{1+\gamma^5}{2} \psi = 0 \Rightarrow$ All mass terms ~~not~~ mixing L w/ R vanish.

$\mathcal{L}_{\text{Dirac}} = (k_c) \bar{\psi}_R \gamma^\mu \partial_\mu \psi_R + (k_c) \bar{\psi}_L \gamma^\mu \partial_\mu \psi_L + m c^2 \bar{\psi}_R \psi_L + m c^2 \bar{\psi}_L \psi_R$

Everything here is in terms of 4-component objects.

$$\mathcal{L}_{\text{Dirac}} = (k_c) \bar{\psi}_R \gamma^\mu \partial_\mu \psi_R + (k_c) \bar{\psi}_L \gamma^\mu \partial_\mu \psi_L + m_c \bar{\psi}_R \psi_L + m_c \bar{\psi}_L \psi_R$$

Now that we have L & R in the game, we need a "doublet" for the $SU(2)_L$ to act on.

Recall for quarks we introduced a triplet $\psi = \begin{pmatrix} \psi_u \\ \psi_d \\ \psi_s \end{pmatrix}$ for $SU(3)$.

We actually pair particles into left-handed doublets: $\chi_L = \begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix}_L, \begin{pmatrix} \nu_\tau \\ \tau \end{pmatrix}_L, \begin{pmatrix} u \\ d \end{pmatrix}_L, \begin{pmatrix} c \\ s \end{pmatrix}_L, \begin{pmatrix} t \\ b \end{pmatrix}_L$

And have right-handed singlets: $e_R, \mu_R, \tau_R, u_R, d_R, c_R, s_R, t_R, b_R$

What about $\nu_{eR}, \nu_{\mu R}, \nu_{\tau R}$? They don't exist! At least not for the massless neutrino story.

For simplicity we will focus on the first generation of leptons, i.e. $\chi_L = \begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, e_R$ and ignore the mass term (since it never plays a role in generating interactions).

$$\mathcal{L}_{\text{Dirac}} = (k_c) \bar{\chi}_L \gamma^\mu \partial_\mu \chi_L + (k_c) \bar{e}_R \gamma^\mu \partial_\mu e_R$$

$$\mathcal{L}_{\text{Dirac}} = (k_L) \bar{\chi}_L \gamma^\mu \partial_\mu \chi_L + (k_R) \bar{e}_R \gamma^\mu \partial_\mu e_R$$

This is invariant under global $SU(2)_L \times U(1)_Y$ where $SU(2)_L: e^{-ig \vec{\sigma} \cdot \vec{\theta}} \chi_L$ $g = \frac{q}{k_L}$
 Isospin Hypercharge $U(1)_Y: e^{-ig' Y \phi} \chi_L, e^{-ig' Y \phi} e_R$

With the factors Y_{k_L} and Y_{k_R} , we allow χ_L and e_R to carry different amounts of the same hypercharge (governed by g'). Each of these can be promoted to a local symmetry of the Lagrangian by the methods we have covered in the preceding lectures.

$$\text{Let: } \partial_\mu \chi_L \rightarrow D_\mu \chi_L = \partial_\mu \chi_L + ig \vec{\sigma} \cdot \vec{W}_\mu \chi_L + ig' Y_{k_L} B_\mu \chi_L$$

3 $SU(2)_L$ gauge fields

So if we adjust g' , then both Y_{k_L} and Y_{k_R} scale accordingly!

$$\partial_\mu e_R \rightarrow D_\mu e_R = \partial_\mu e_R + ig' Y_{k_R} B_\mu e_R$$

Where: $\vec{\sigma} \cdot \vec{W}'_\mu = e^{-ig \vec{\sigma} \cdot \vec{\theta}} \vec{\sigma} \cdot \vec{W}_\mu e^{ig \vec{\sigma} \cdot \vec{\theta}} + \frac{i}{g} \partial_\mu (e^{-ig \vec{\sigma} \cdot \vec{\theta}}) e^{ig \vec{\sigma} \cdot \vec{\theta}}$ } Gauge field transformation rules.
 $B'_\mu = B_\mu + \partial_\mu \phi$

Then for each gauge field we introduce a kinetic term using $F_{\mu\nu} = -\frac{i}{g} [D_\mu, D_\nu]$ and again this would give the usual $\begin{cases} F_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu & \text{for the } U(1)_Y \\ F_{\mu\nu} = \partial_\mu W_\nu - \partial_\nu W_\mu - g \epsilon^{abc} W_\mu^a W_\nu^b & \text{for } SU(2)_L \end{cases}$

The latter will include interactions between the $SU(2)_L$ gauge bosons (just like the gluons of QCD). $SU(2)_L$ Lie algebra structure constants

The $SU(2)_L$ interactions have a particularly interesting structure. We expect 3 gauge bosons $W_\mu^1, W_\mu^2, W_\mu^3$ which act on the left-handed doublets, e.g. $\chi_L = \begin{pmatrix} \nu_e \\ e \end{pmatrix}_L$.

Using the machinery of good old spin-1/2 from QM (that involved $SU(2)$ as well!) we can think of this in terms of:

$$\begin{aligned} W_\mu^3 &\sim i\sigma^3 = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix} \Rightarrow W_\mu^3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} = +\frac{1}{2} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad W_\mu^3 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = -\frac{1}{2} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ W_\mu^+ &\sim \frac{1}{\sqrt{2}} (-i\sigma^1 - \sigma^2) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \Rightarrow W_\mu^+ \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 0 \quad W_\mu^+ \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ W_\mu^- &\sim \frac{1}{\sqrt{2}} (i\sigma^1 - \sigma^2) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \Rightarrow W_\mu^- \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad W_\mu^- \begin{pmatrix} 0 \\ 1 \end{pmatrix} = 0 \end{aligned}$$

↑ These contain the action in flavor-space.

But notice that W_μ^+ (W_μ^-) changes $e \rightarrow \nu_e$ ($\nu_e \rightarrow e$) respectively, so clearly the \pm here must be connected to electric charge. These gauge bosons mediate interactions which change the electric charge of the matter involved.

The W_μ^3 on the other hand does not change the particle type, hence does not change the electric charge, i.e. it is neutral.

There are 2 big problems:

i) We know that the weak gauge bosons are massive, and we already know that the Proca mass term $(\frac{m}{\hbar})^2 W_\mu^\alpha W^{\mu\alpha}$ is not gauge invariant.

ii) Recall that to have mass terms for spinors requires both the L and R parts of ψ to combine, e.g. $m\psi_L^\dagger \psi_R$. However we have just constructed a gauge theory where the L and R parts transform differently. There is no way we can expect $m\psi_L^\dagger \psi_R$ to be gauge invariant!

Both of these problems will be solved w/ the Higgs mechanism for mass generation. A crucial part of this process is the breaking of $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$.

We will have much more to say about how such a symmetry can be broken, but for now we will just highlight the implication for the electroweak interactions.

$SU(2)_L \times U(1)_Y$ has 4 generators $W_\mu^3, W_\mu^\pm, B_\mu$. After symmetry breaking to $U(1)_{EM}$ we only expect one symmetry generator to survive. Which one is it?

You might have thought it would be B_μ , then $U(1)_Y \rightarrow U(1)_{EM}$, but that is not the case.

In actuality, the B_μ "mixes" with the neutral W_μ^3 from $SU(2)_L$. We can form 2 orthogonal states:

$$\begin{aligned} A_\mu &= B_\mu \cos\theta_w + W_\mu^3 \sin\theta_w \Rightarrow \text{The photon of } U(1)_{EM} \\ Z_\mu &= -B_\mu \sin\theta_w + W_\mu^3 \cos\theta_w \Rightarrow \text{The massive neutral } Z^0 \text{ boson of the weak interactions.} \end{aligned}$$

\uparrow
 $\theta_w = \text{Weinberg mixing angle}$

So it should be clear that we cannot identify just $U(1)_{EM}$ with the $U(1)$ factor in $SU(2)_L \times U(1)_Y$!

Going back to the original unified gauge group $SU(2)_L \times U(1)_Y$, it wouldn't make much sense to call this a "unified" group if the $SU(2)_L$ and $U(1)_Y$ factors had completely independent couplings g and g' . It turns out that they are in fact related. However since we experience the broken version of this theory, it is more useful to know how the couplings to W^\pm, Z^0 and γ are related.

Turns out: $g \sin\theta_w = g' \cos\theta_w = g_\gamma$ and $g = g_{W^\pm}$ while $g_Z = \frac{g_\gamma}{\sin\theta_w \cos\theta_w}$